

Equivariant autoequivalences for finite group actions

David Ploog

ABSTRACT The familiar Fourier-Mukai technique can be extended to an equivariant setting where a finite group G acts on a smooth projective variety X . In this paper we compare the group of invariant autoequivalences $\text{Aut}(\text{D}^b(X))^G$ with the group of autoequivalences of $\text{D}^G(X)$. We apply this method in three cases: Hilbert schemes on K3 surfaces, Kummer surfaces and canonical quotients.

1 Introduction and Setup

It often proves useful to consider analogues of classical settings, adding the presence of a group action. Instances of this in algebraic geometry are e.g. equivariant intersection theory or the McKay type theorems. There already is a theory for derived categories of varieties with actions by algebraic groups [2]. In this article, we study the behaviour of automorphism groups of such derived categories in the case when the group is finite. This paper grew out of my Ph.D. thesis [16]. I would like to thank Georg Hein, Daniel Huybrechts, Manfred Lehn, and Richard Thomas for their help and valuable suggestions.

We always work with varieties over $\mathbb{C}$. A kernel $P \in \text{D}^b(X \times Y)$ gives rise to a Fourier-Mukai transform which we denote by $\text{FM}_P : \text{D}^b(X) \rightarrow \text{D}^b(Y)$. We also introduce special notation for composition of such transforms: Let us write $\text{FM}_Q \circ \text{FM}_P = \text{FM}_{Q \star P} : \text{D}^b(X) \rightarrow \text{D}^b(Z)$ for $P \in \text{D}^b(X \times Y)$ and $Q \in \text{D}^b(Y \times Z)$, i.e. $Q \star P = \mathbb{R}p_{XZ*}(p_{XY}^*P \otimes^{\mathbb{L}} p_{YZ}^*Q)$. This works for morphisms as well: $f : P \rightarrow P'$ and $g : Q \rightarrow Q'$ give rise to $g \star f : Q \star P \rightarrow Q' \star P'$.

1.1 Linearisations and $\text{D}^G(X)$

Let X be a smooth projective variety on which a finite group G acts. A G -linearisation of a sheaf E on X is given by isomorphisms $\lambda_g : E \xrightarrow{\sim} g^*E$ for all $g \in G$ satisfying $\lambda_1 = \text{id}_E$ and $\lambda_{gh} = h^*\lambda_g \circ \lambda_h$. A morphism $f : (E_1, \lambda_1) \rightarrow (E_2, \lambda_2)$ is G -invariant, if $f = g \cdot f := \lambda_{2,g}^{-1} \circ g^*f \circ \lambda_{1,g}$ for all $g \in G$. The category of G -linearised coherent sheaves on X with G -invariant morphisms is denoted by $\text{Coh}^G(X)$; note that it is abelian and contains enough injectives, see [3]. Put $\text{D}^G(X) := \text{D}^b(\text{Coh}^G(X))$ for its derived category.

There is an equivalent point of view on $\text{D}^G(X)$: let $\mathcal{T}$ be the category consisting of G -linearised objects of $\text{D}^b(X)$, i.e. complexes $E^\bullet \in \text{D}^b(X)$ together with isomorphisms $\lambda_g : E^\bullet \xrightarrow{\sim} g^*E^\bullet$ in $\text{D}^b(X)$ satisfying the same co-cycle condition as above. The canonical functor $\text{D}^G(X) \rightarrow \mathcal{T}$ is fully faithful in view of $\text{Hom}_{\text{D}^G(X)}((E_1, \lambda_1)^\bullet, (E_2, \lambda_2)^\bullet) = \text{Hom}_{\text{D}^b(X)}(E_1^\bullet, E_2^\bullet)^G$ for objects $(E_1, \lambda_1)^\bullet, (E_2, \lambda_2)^\bullet \in \text{D}^G(X)$. To show that the functor is essentially surjective, take $(E^\bullet, \lambda) \in \mathcal{T}$. Choosing an injective bounded resolution $E^\bullet \xrightarrow{\sim} I^\bullet$, we can assume that λ corresponds to genuine complex maps $\tilde{\lambda}_g : I^\bullet \xrightarrow{\sim} g^*I^\bullet$. Hence $(I, \tilde{\lambda})^\bullet$ is a complex with a linearisation of each sheaf, and using $\text{D}^b(X) \cong \text{D}_{\text{Coh}(X)}^b(\text{Qcoh}(X)) \ni (I, \tilde{\lambda})^\bullet$, we get $\mathcal{T} \cong \text{D}^b(X)$.

1.2 Equivariant push-forwards and Fourier-Mukai transforms

Now consider two such varieties with finite group actions, (X, G) and (X', G') . A map between them is given by a pair of morphisms $\Phi : X \rightarrow X'$ and $\varphi : G \rightarrow G'$ such that $\Phi \circ g = \varphi(g) \circ \Phi$ for all $g \in G$. Then, we have the pull-back $\Phi^* : \text{Coh}^{G'}(X') \rightarrow \text{Coh}^G(X)$ (and its derived functor $\mathbb{L}\Phi^* : \text{D}^{G'}(X') \rightarrow \text{D}^G(X)$) which just means equipping the usual pull-back Φ^*E' with the G_1 -linearisation $\Phi^*\lambda'_{\varphi(g)} : \phi^*E' \simeq \Phi^*\varphi(g)^*E' = g^*\Phi^*E'$.

Suppose that φ is surjective and put $K := \ker(\varphi)$. Then, there is also an equivariant push-forward defined for $(E, \lambda) \in \text{Coh}^G(X)$ in the following way: the usual push-forward Φ_*E is canonically G -linearised since φ is surjective. Now K acts trivially on X' , thus is it possible to take K -invariants. Then the subsheaf $\Phi_*^K(E, \lambda) := [\Phi_*E]^K \subset \Phi_*E$ is still G' -linearised and $\mathbb{R}\Phi_*^K : \text{D}^G(X) \rightarrow \text{D}^{G'}(X')$ is the correct push-forward.

For objects $(E, \lambda) \in \text{Coh}^G(X)$ and $(E', \lambda') \in \text{Coh}^{G'}(X')$ and a G -invariant morphism $\Phi^*E' \rightarrow E$, the adjoint morphism $E' \rightarrow \Phi_*E$ has image in Φ_*^KE . Hence, the functors $\Phi^* : \text{Coh}^{G'}(X') \rightarrow \text{Coh}^G(X)$ and $\Phi_*^K : \text{Coh}^G(X) \rightarrow \text{Coh}^{G'}(X')$ are adjoint; analogously for $\mathbb{L}\Phi^*$ and $\mathbb{R}\Phi_*^K$.

As a consequence, the usual Fourier-Mukai calculus extends to the equivariant setting if we use these functors (the tensor product of two linearised objects is obviously again linearised). Explicitly, for an object $(P, \rho) \in \text{D}^{G \times G'}(X \times X')$ we get a functor

$$\text{FM}_{(P, \rho)} : \text{D}^G(X) \rightarrow \text{D}^{G'}(X'), \quad (E, \lambda) \mapsto \mathbb{R}p_{X'}^G(P \otimes^{\mathbb{L}} p_X^*(E))$$

with the projections $p_{X'} : X \times X' \rightarrow X'$ and $p_X : X \times X' \rightarrow X$ (and similar projections on the group level).

1.3 Inflation and restriction

There is an obvious forgetful functor $for : \text{D}^G(X) \rightarrow \text{D}^b(X)$. In the other direction, we have the inflation functor $inf : \text{D}^b(X) \rightarrow \text{D}^G(X)$ with $inf(E) := \bigoplus_{g \in G} g^*E$ and the G -linearisation comes from permuting the summands¹. A generalisation of inf to the case of a subgroup $H \subset G$ is given by

$$inf_H^G : \text{D}^H(X) \rightarrow \text{D}^G(X), \quad (E, \lambda) \mapsto \bigoplus_{[g] \in H \backslash G} g^*E$$

and the G -linearisation of the sum is a natural combination of λ and permutations.

See Bernstein/Lunts [2] for generalisations of $\text{D}^G(X)$ and inf_H^G to the case of algebraic groups (neither of which is straight forward).

¹ In a similar vein, every symmetric polynomial $\sigma \in \mathbb{Z}[x_1, \dots, x_n]^{S_n}$ (where $n := \#G$) gives rise to a functor $inf_\sigma : \text{D}^b(X) \rightarrow \text{D}^G(X)$. For example, $inf_{x_1 + \dots + x_n} = inf$ and $inf_{x_1 \dots x_n}(E) = \bigotimes_{g \in G} g^*E$. As an application, for an ample line bundle L on X we get $inf_{x_1 \dots x_n}(L)$, an ample G -linearised line bundle.

1.4 Invariant vs linearised objects

Obviously, a G -linearised object has to be G -invariant, i.e. fixed by all pull-backs g^* . It is a difficult question under which conditions the other direction is true. For us the following fact ([16, Lemma 3.4]) will suffice.

Lemma 1. *Let $E \in D^b(X)$ be simple and G -invariant. Then there is a group cohomology class $[E] \in H^2(G, \mathbb{C}^*)$ such that E is G -linearisable if and only if $[E] = 0$. Furthermore, if $[E] = 0$, then the set of G -linearisations of E is canonically a $\hat{G}$ -torsor.*

Proof. Note that the G -action on $\text{Aut}(E) = \mathbb{C}^*$ is trivial. There are isomorphisms $\mu_g : E \xrightarrow{\sim} g^*E$ for all $g \in G$. As E is simple, we can define units $c_{g,h} \in \mathbb{C}^*$ by $\mu_{gh} = h^*\mu_g \circ \mu_h \cdot c_{g,h}$. It is a straightforward check that the map $c : G^2 \rightarrow \mathbb{C}^*$ is a 2-cocycle of G with values in $\mathbb{C}^*$, i.e. $c \in Z^2(G, \mathbb{C}^*)$. Replacing the isomorphisms μ_g with some other μ'_g yields the map $e : G \rightarrow \mathbb{C}^*$ such that $\mu'_g = \mu_g \cdot e_g$. The two cocycles $c, c' : G^2 \rightarrow \mathbb{C}^*$ derived from μ and μ' differ by the boundary coming from e by another easy computation. Hence, $c/c' = d(e)$ and thus $c = c' \in H^2(G, \mathbb{C}^*)$. Thus the G -invariant object E gives rise to a unique class $[E] := c \in H^2(G, \mathbb{C}^*)$. In these terms, E is G -linearisable if and only if $c \equiv 1$, i.e. $[E]$ vanishes.

For the second statement, we write $\hat{G} := \text{Hom}(G, \mathbb{C}^*)$ for the group of characters and $\text{Lin}_G(E)$ for the set of non-isomorphic G -linearisations of E . Consider the $\hat{G}$ -action $\hat{G} \times \text{Lin}_G(E) \rightarrow \text{Lin}_G(E)$, $(\chi, \lambda) \mapsto \chi \cdot \lambda$ on $\text{Lin}_G(E)$. First take $\chi \in \hat{G}$ and $\lambda \in \text{Lin}_G(E)$ such that $\chi \cdot \lambda = \lambda$. Then, there is an isomorphism $f : (E, \lambda) \xrightarrow{\sim} (E, \chi \cdot \lambda)$ which in turn immediately implies $\chi = 1$ using $f \in \text{Aut}(E) = \mathbb{C}^*$. Thus, the action is effective. Now take two elements $\lambda, \lambda' \in \text{Lin}_G(E)$ and consider $\lambda_g^{-1} \circ \lambda'_g : E \xrightarrow{\sim} g^*E \xrightarrow{\sim} E$. As E is simple, we have $\lambda_g^{-1} \circ \lambda'_g = \chi(g) \cdot \text{id}_E$. It follows from the cocycle condition for linearisations that χ is multiplicative, i.e. $\chi \in \hat{G}$. In other words, $\lambda' = \chi \cdot \lambda$ and the action is also transitive. Altogether $\hat{G}$ acts simply transitive on $\text{Lin}_G(E)$. $\square$

For our use of group cohomology, refer for example [17]. The second cohomology $H^2(G, \mathbb{C}^*)$ of the finite group G acting trivially (on an algebraically closed field of characteristic 0) is also known as the *Schur multiplier* of G (refer [8, §25]). Two relevant facts about it are: $H^2(G, \mathbb{C}^*)$ is a finite abelian group; its exponent is a divisor of $\#G$. Examples are given by $H^2((\mathbb{Z}/n\mathbb{Z})^k, \mathbb{C}^*) = (\mathbb{Z}/n\mathbb{Z})^{k(k-1)/2}$ for copies of a cyclic group; $H^2(D_{2n}, \mathbb{C}^*) = \mathbb{Z}/2\mathbb{Z}$ and $H^2(D_{2n+1}, \mathbb{C}^*) = 0$ for the dihedral groups with $n > 1$; and $H^2(S_n, \mathbb{C}^*) = \mathbb{Z}/2\mathbb{Z}$ for the symmetric groups with $n > 3$.

Note that a group with vanishing Schur multiplier has the following property: every simple G -invariant object of $D^G(X)$ is G -linearisable, no matter how G acts on X .

Remark 2. The condition that E be simple in the Lemma is important. Consider an abelian surface A with the action of $G = \mathbb{Z}/2\mathbb{Z} = \{\pm \text{id}_A\}$. Then the sheaf $E := k(a) \oplus k(-a) = \text{inf}(k(a))$ is G -invariant but not simple. Yet it is uniquely $\mathbb{Z}/2\mathbb{Z}$ -linearisable as an easy computation shows [16, Example 3.9] (in contrast to G -invariant simple sheaves, which have precisely two non-isomorphic G -linearisations according to $H^2(\mathbb{Z}/2\mathbb{Z}, \mathbb{C}^*) = 0$). This behaviour is expected from geometry: by derived McKay correspondence ([3]) one has

$D^G(A) \cong D^b(X)$, where X is the Kummer surface of A , a crepant resolution $\psi : X \rightarrow A/G$. Under this equivalence, skyscraper sheaves of points $x \in X$ outside of exceptional fibres of ψ are mapped to $k(\psi(x)) \oplus k(-\psi(x))$.

Example 3. If as before G acts on X , then the canonical sheaf ω_X is simple (as it is a line bundle) and G -invariant (because it is functorial). Due to this functoriality, it is actually G -linearisable: the morphism $g : X \rightarrow X$ induces a morphism of cotangent bundles $g_* : g^* \Omega_X \rightarrow \Omega_X$. Going to determinants and using adjunction yields the desired isomorphisms $\lambda_g := \det(g_*^{-1}) : \omega_X \xrightarrow{\sim} g^* \omega_X$.

2 Groups of autoequivalences

We are interested in comparing the automorphism group $\text{Aut}(D^G(X))$ with the group $\text{Aut}(D^b(X))^G := \{F \in \text{Aut}(D^b(X)) : g^* \circ F = F \circ g^* \forall g \in G\}$. It turns out that a useful intermediate step is to look at Fourier-Mukai equivalences on $D^b(X)$ which are diagonally G -linearised.

To make this precise, consider a Fourier-Mukai transform $\text{FM}_P : D^b(X) \rightarrow D^b(X')$. Suppose that G acts on both X and X' . Then we have the diagonal action $G \times X \times X' \rightarrow X \times X'$, $g \cdot (x, x') := (gx, gx')$ which we sometimes (especially in the case $X = X'$) for emphasis call the G_Δ -action of G on $X \times X'$. Now we are in a position to study objects $(P, \rho) \in D^{G_\Delta}(X \times X')$ which give Fourier-Mukai equivalences $\text{FM}_P : D^b(X) \xrightarrow{\sim} D^b(X')$. In other words, these are ordinary kernels for equivalences $D^b(X) \xrightarrow{\sim} D^b(X')$ which additionally have been equipped with a G_Δ -linearisation.

Not every kernel in $P \in D^b(X \times X')$ has the latter property. A necessary condition is that P must be G_Δ -invariant, i.e. $(g, g)^* P \cong P$ for all $g \in G$, or, equivalently, $g^* \circ \text{FM}_P = \text{FM}_P \circ g^*$. Now we apply the following general fact:

Lemma 4. *If $P \in D^b(X \times Y)$ is the Fourier-Mukai kernel of an equivalence $\text{FM}_P : D^b(X) \xrightarrow{\sim} D^b(Y)$ then P is simple, i.e. $\text{Hom}_{D^b(X \times Y)}(P, P) = \mathbb{C}$.*

Proof. Fix $f \in \text{Hom}_{D^b(X \times Y)}(P, P)$ and let Q be a quasi-inverse kernel for FM_P , i.e. $P \star Q \cong \mathcal{O}_{\Delta_Y}$. By $\text{Hom}_{D^b(Y \times Y)}(\mathcal{O}_{\Delta_Y}, \mathcal{O}_{\Delta_Y}) = \text{Hom}_{Y \times Y}(\mathcal{O}_{\Delta_Y}, \mathcal{O}_{\Delta_Y}) = \mathbb{C}$ we have $f \star \text{id}_Q = c \cdot \text{id}_{\mathcal{O}_{\Delta_Y}}$ for a $c \in \mathbb{C}$. Composing again with $\text{id}_P : P \rightarrow P$ gives $f = f \star \text{id}_{\mathcal{O}_{\Delta_X}} = f \star (\text{id}_Q \star \text{id}_P) = (f \star \text{id}_Q) \star \text{id}_P = (c \cdot \text{id}_{\mathcal{O}_{\Delta_Y}}) \star \text{id}_P = c \cdot \text{id}_P$. $\square$

Combining Lemmas 1 and 4, we see that G_Δ -invariant kernels for equivalences are G_Δ -linearisable, provided that $H^2(G, \mathbb{C}^*) = 0$ (or more generally, if the obstruction class in $H^2(G, \mathbb{C}^*)$ vanishes).

Now suppose we have an arbitrary object $(P, \rho) \in D^{G_\Delta}(X \times X')$ and the accompanying functor $\text{FM}_P : D^b(X) \rightarrow D^b(X')$. The general device of inflation allows us to produce the following equivariant Fourier-Mukai transform from (P, ρ) :

$$\text{FM}_{(P, \rho)}^G := \text{FM}_{\inf_{G_\Delta}^{G^2}(P, \rho)} : D^G(X) \rightarrow D^G(X').$$

For brevity, we set $G \cdot P := \inf_{G_\Delta}^{G^2}(P, \rho) \in D^{G^2}(X \times X')$ and call it *the inflation of (P, ρ)* . The following lemma states the main properties of this assignment.

Lemma 5. *Let X, X', X'' be smooth projective varieties with G -actions and let $(P, \rho) \in D^{G_\Delta}(X \times X')$ and $(P', \rho') \in D^{G_\Delta}(X' \times X'')$.*

(1) $\text{FM}_{(\mathcal{O}_{\Delta_X}, \text{can})}^G \cong \text{id} : D^G(X) \rightarrow D^G(X)$.

(2) For $(P, \rho) \in D^{G\Delta}(X \times X')$ there is a commutative diagram

$$\begin{array}{ccc} D^G(X) & \xrightarrow{\text{FM}_{(P,\rho)}^G} & D^G(X') \\ \downarrow \text{for} & & \downarrow \text{for} \\ D^b(X) & \xrightarrow{\text{FM}_P} & D^b(X') \end{array}$$

(3) $\text{FM}_{(P',\rho')}^G \circ \text{FM}_{(P,\rho)}^G \cong \text{FM}_{(P' \star P, \rho' \star \rho)}^G$ where $(\rho' \star \rho)_g := \rho'_g \star \rho_g$.

(4) FM_P fully faithful $\implies \text{FM}_{(P,\rho)}^G$ fully faithful.

(5) FM_P equivalence $\implies \text{FM}_{(P,\rho)}^G$ equivalence.

Proof. (1) The structure sheaf $\mathcal{O}_\Delta$ of the diagonal $\Delta \subset X \times X$ has a canonical G_Δ -linearisation, as $(g, g)^* \mathcal{O}_\Delta = \mathcal{O}_\Delta$. The inflation of $\mathcal{O}_\Delta$ is $G \cdot \mathcal{O}_\Delta = \bigoplus_{g \in G} \mathcal{O}_{(g,1)\Delta}$, and its G^2 -linearisation is given by the permutation of summands via $G \rightarrow G, g \mapsto kgh^{-1}$.

Using this one can check by hand that $\text{FM}_{(\mathcal{O}_\Delta, \text{can})}^G$ maps any $(E, \lambda) \in D^G(X)$ to itself (see [16, Example 3.14]).

(2) Take any object $(E, \lambda) \in D^G(Y)$. Then, we have by definition of equivariant Fourier-Mukai transforms $\text{FM}_P^G : D^G(Y) \rightarrow D^G(Y)$, $(E, \lambda) \mapsto [\mathbb{R}p_{2*}(G \cdot P \otimes^{\mathbb{L}} p_1^* E)]^{G \times 1}$. The $G \times 1$ -linearisation of $G \cdot P$ is given by permutations (the G_Δ -linearisation of P does not enter). Since $\mathbb{R}p_{2*}((g, 1)^* P \otimes^{\mathbb{L}} p_1^* E) = \mathbb{R}p_{2*}(g, 1)^*(P \otimes^{\mathbb{L}} p_1^* g^{-1*} E) = \mathbb{R}p_{2*}(P \otimes^{\mathbb{L}} p_1^* E)$, we see that $\mathbb{R}p_{2*}(G \cdot P \otimes^{\mathbb{L}} p_1^* E) \cong \bigoplus_G \mathbb{R}p_{2*}(P \otimes^{\mathbb{L}} p_1^* E)$ and $G \times 1$ acts with permutation matrices where the 1's are replaced by $p_1^* \lambda_g$'s. Taking $G \times 1$ -invariants singles out a subobject of this sum isomorphic to one summand.

A morphism $f : E_1 \rightarrow E_2$ in $D^G(X)$ is likewise first taken to a G -fold direct sum. The final taking of $G \times 1$ -invariants then leaves one copy of $\text{FM}_P(f)$.

(3) The composite $\text{FM}_{G \cdot P'} \circ \text{FM}_{G \cdot P}$ has the kernel

$$\begin{aligned} (G \cdot P') \star (G \cdot P) &= [\mathbb{R}p_{13*}(p_{12}^*(G \cdot P) \otimes^{\mathbb{L}} p_{23}^*(G \cdot P'))]^{1 \times G \times 1} \\ &= [\mathbb{R}p_{13*}(p_{12}^* \bigoplus_{g \in G} (g, 1)^* P \otimes^{\mathbb{L}} p_{23}^* \bigoplus_{h \in G} (h, 1)^* P')]^{1 \times G \times 1} \\ &\cong_{\rho'} [\mathbb{R}p_{13*}(p_{12}^* \bigoplus_{g \in G} (g, 1)^* P \otimes^{\mathbb{L}} p_{23}^* \bigoplus_{h \in G} (1, h^{-1})^* P')]^{1 \times G \times 1} \\ &= [\bigoplus_{g, h \in G} \mathbb{R}p_{13*}(g, 1, h^{-1})^*(p_{12}^* P \otimes^{\mathbb{L}} p_{23}^* P')]^{1 \times G \times 1} \\ &= [\bigoplus_{g, h \in G} (g, h^{-1})^* \mathbb{R}p_{13*}(p_{12}^* P \otimes^{\mathbb{L}} p_{23}^* P')]^{1 \times G \times 1}. \end{aligned}$$

Now note that $(1, c, 1) \in 1 \times G \times 1$ acts on $(G \cdot P') \star (G \cdot P)$ via permutations (inverse multiplications from left) and ρ on P , and $(1, c, 1)$ acts purely by permutations (which are multiplications from right) on P' . Plugging this into the last equation, we find that after taking invariants we end up with $\bigoplus_{d \in G} (d^{-1}, d)^* \mathbb{R}p_{13*}(p_{12}^* P \otimes^{\mathbb{L}} p_{23}^* P')$. Since the (d^{-1}, d) 's give all classes in $G_\Delta \setminus G^2$, we find that $(G \cdot P') \star (G \cdot P) \cong G \cdot (P' \star P)$.

(4) Fix two objects $(E_1, \lambda_1), (E_2, \lambda_2) \in D^G(X)$. The injectivity of the natural map $\text{Hom}_{D^G(X)}((E_1, \lambda), (E_2, \lambda_2)) \rightarrow \text{Hom}_{D^G(X')}(\text{FM}_P^G(E_1, \lambda_1), \text{FM}_P^G(E_2, \lambda_2))$ is

a consequence of two facts: $\mathrm{Hom}_{\mathrm{D}^G(X)}(\cdot, \cdot) = \mathrm{Hom}_{\mathrm{D}^b(X)}(\cdot, \cdot)^G \subset \mathrm{Hom}_{\mathrm{D}^b(X)}(\cdot, \cdot)$ by the definition of morphisms in $\mathrm{D}^G(X)$ on one hand and $\mathrm{Hom}_{\mathrm{D}^b(X)}(E_1, E_2) \cong \mathrm{Hom}_{\mathrm{D}^b(X')}(\mathrm{FM}_P(E_1), \mathrm{FM}_P(E_2))$ by hypothesis.

Similarly, the surjectivity uses the same facts. One just replaces the embedding $\mathrm{Hom}_{\mathrm{D}^b(X)}(\cdot, \cdot)^G \subset \mathrm{Hom}_{\mathrm{D}^b(X)}(\cdot, \cdot)$ with the averaging projection (Reynolds operator) $\theta : \mathrm{Hom}_{\mathrm{D}^b(X)}(E_1, E_2) \rightarrow \mathrm{Hom}_{\mathrm{D}^b(X)}(E_1, E_2)^G$ given by $\theta(f) := \frac{1}{\#G} \sum_{g \in G} \lambda_{2,g}^{-1} \circ g^* f \circ \lambda_{1,g}$.

(5) follows from (3) and (1): let $\mathrm{FM}_P : \mathrm{D}^b(X) \xrightarrow{\sim} \mathrm{D}^b(X')$ be an equivalence. Then, $Q = \mathbb{R}\mathrm{Hom}(P, \mathcal{O}_{X \times X'}) \otimes p_X^* \omega_X[\dim(X)]$ is the Fourier-Mukai kernel of a quasi-inverse for FM_P . As the canonical bundle ω_X is G -linearisable (see Example 3), Q inherits a G_Δ -linearisation from those of P and $p_X^* \omega_X$. We have $(G \cdot P) \star (G \cdot Q) = G \cdot (P \star Q) = G \cdot \mathcal{O}_\Delta$. There are precisely $\#\hat{G}$ different G -linearisations for ω_X as well as $\#\hat{G}$ different G_Δ -linearisations for $\mathcal{O}_\Delta$. Thus, exactly one choice of G -linearisation for ω_X will equip the composition $P \star Q = \mathcal{O}_\Delta$ with the canonical G_Δ -linearisation. But then (1) shows that this is the kernel of the identity on $\mathrm{D}^G(X')$. Hence, $G \cdot P$ is an equivalence kernel as was P . $\square$

Let us consider the following automorphism groups:

$$\begin{aligned} \mathrm{Aut}(\mathrm{D}^b(X))^G &= \{\mathrm{FM}_P \in \mathrm{Aut}(\mathrm{D}^b(X)) : (g, g)^* P \cong P \ \forall g \in G\} \\ \mathrm{Aut}(\mathrm{D}^G(X)) &= \{\mathrm{FM}_{\tilde{P}} : \mathrm{D}^G(X) \xrightarrow{\sim} \mathrm{D}^G(X) : \tilde{P} \in \mathrm{D}^{G^2}(X \times X)\} \\ \mathrm{Aut}^{G_\Delta}(\mathrm{D}^b(X)) &:= \{(P, \rho) \in \mathrm{D}^{G_\Delta}(X \times X) : \mathrm{FM}_P \in \mathrm{Aut}(\mathrm{D}^b(X))\}. \end{aligned}$$

The first identity uses the action $G \times \mathrm{Aut}(\mathrm{D}^b(X)) \rightarrow \mathrm{Aut}(\mathrm{D}^b(X))$ given by $g \cdot F := (g^{-1})^* \circ F \circ g^*$ and the formula $\mathrm{FM}_{(g,g)^* P} = g^* \circ \mathrm{FM}_P \circ (g^{-1})^*$ and finally Orlov's result on the existence of Fourier-Mukai kernels [14, Theorem 2.2]. This has been extended by Kawamata to smooth stacks associated to normal projective varieties with quotient singularities [11]. In view of $\mathrm{Coh}([X/G]) \cong \mathrm{Coh}^G(X)$, this implies the second relation. Finally we have to turn $\mathrm{Aut}^{G_\Delta}(\mathrm{D}^b(X))$ into a group. This is done by Lemma 5: (3) settles the composition, (1) the neutral element and (5) the inverses.

The following theorem [16, Proposition 3.17] is an attempt to compare these groups.

Theorem 6. *Let the finite group G act on a smooth projective variety X .*

(1) *The construction of inflation gives a group homomorphism inf which fits in the following exact sequence, where $Z(G) \subset G$ is the centre of G :*

$$\begin{array}{ccccccc} 0 & \longrightarrow & Z(G) & \longrightarrow & \mathrm{Aut}^{G_\Delta}(\mathrm{D}^b(X)) & \xrightarrow{\mathrm{inf}} & \mathrm{Aut}(\mathrm{D}^G(X)) \\ & & & & (P, \rho) & \longmapsto & \mathrm{FM}_{(P, \rho)}^G \end{array}$$

(2) *Forgetting the G_Δ -linearisation gives a group homomorphism for which sits in the following exact sequence; here $G_{\mathrm{ab}} := G/[G, G] \cong \mathrm{Hom}(G, \mathbb{C})^* = H^1(G, \mathbb{C}^*)$ is the abelianisation:*

$$\begin{array}{ccccccc} 0 & \longrightarrow & G_{\mathrm{ab}} & \longrightarrow & \mathrm{Aut}^{G_\Delta}(\mathrm{D}^b(X)) & \xrightarrow{\mathrm{for}} & \mathrm{Aut}(\mathrm{D}^b(X))^G \longrightarrow H^2(G, \mathbb{C}^*) \\ & & & & (P, \rho) & \longmapsto & \mathrm{FM}_P \longmapsto [P] \end{array}$$

Proof. (1) It follows from Lemma 5 that inf is a group homomorphism. The kernel $\ker(\text{inf})$ consists of $(P, \rho) \in \text{D}^{G_\Delta}(X^2)$ giving equivalences such that $G \cdot P \cong G \cdot \mathcal{O}_\Delta$. Obviously, this forces P to be a sheaf of type $P \cong (g, 1)^* \mathcal{O}_\Delta$ for some $g \in G$. Now $(g, 1)^* \mathcal{O}_\Delta$ is G_Δ -invariant if and only if $g \in Z(G)$ as $(h, h)^*(g, 1)^* \mathcal{O}_\Delta = (gh, h)^* \mathcal{O}_\Delta \cong (gh, h)^*(h^{-1}, h^{-1})^* \mathcal{O}_\Delta = (h^{-1}gh, 1)^* \mathcal{O}_\Delta$. This in turn implies that the isomorphism $(g, 1) : X^2 \xrightarrow{\sim} X^2$ is a G_Δ -map. In particular, $P \cong (g, 1)^* \mathcal{O}_\Delta$ gets the pulled back G_Δ -linearisation. Giving $\mathcal{O}_\Delta$ a G_Δ -linearisation $\lambda \in \hat{G}$ different from the canonical one yields $G \cdot (\mathcal{O}_\Delta, \lambda) \not\cong G \cdot (\mathcal{O}_\Delta, \text{can})$; this follows for example from the uniqueness of Fourier-Mukai kernels. Both facts together imply $\ker(\text{inf}) \cong Z(G)$.

(2) It is obvious from the definition of $\text{Aut}^{G_\Delta}(\text{D}^b(X))$ that for is a group homomorphism. The kernel of for corresponds to the G_Δ -linearisations on $\mathcal{O}_\Delta$. From Lemma 1, we see that they form a group isomorphic to $\hat{G} = \text{Hom}(G, \mathbb{C}^*) \cong G_{\text{ab}}$. Note also $\text{Hom}(G, \mathbb{C}^*) = H^1(G, \mathbb{C}^*)$ as G acts trivial on $\mathbb{C}^*$. Given $\text{FM}_P \in \text{Aut}(\text{D}^b(X))^{G_\Delta}$, we know from Lemma 4 that its Fourier-Mukai kernel P is simple. Furthermore, it is G_Δ -invariant by assumption, so that the map $\text{FM}_P \mapsto [P]$ is defined as in Lemma 1. To see that it is a group homomorphism, take two G_Δ -invariant kernels $P, Q \in \text{D}^b(X^2)$. Choose isomorphisms $\lambda_g : P \xrightarrow{\sim} (g, g)^* P$ and $\mu_g : Q \xrightarrow{\sim} (g, g)^* Q$ for all $g \in G$. Then we have $\lambda_{gh} = (h, h)^* \lambda_g \circ \lambda_h \cdot [P]_{g, h}$ and likewise for Q, μ . Furthermore, the composition of λ_g and μ_g gives an isomorphism $\mu_g \star \lambda_g : Q \star P \xrightarrow{\sim} ((g, g)^* Q) \star ((g, g)^* P)$ and the latter term is canonically isomorphic to $(g, g)^*(Q \star P)$. Then $\mu_{g, h} \star \lambda_{gh} = (h, h)^*(\mu_g \star \lambda_g) \circ (\mu_h \star \lambda_h) \cdot [Q \star P]_{g, h}$. The formula $(B \circ A) \star (D \circ C) \cong (B \star D) \circ (A \star C)$ for morphisms $P'' \xrightarrow{B} P' \xrightarrow{A} P$ and $Q'' \xrightarrow{D} Q' \xrightarrow{C} Q$ implies $[Q \star P]_{g, h} = [Q]_{g, h} \cdot [P]_{g, h}$. Now it is obvious that $[\cdot] \circ \text{for} = 0$ and finally $\text{FM}_P \in \text{Aut}(\text{D}^b(X))^{G_\Delta}$ with $[P] = 0$ implies that P is G_Δ -linearisable by Lemma 1. $\square$

3 Applications

The applications will make use of the derived McKay correspondence: assume that X is a smooth (quasi-)projective variety with an action of the finite group G . Then there is a G -Hilbert scheme $G\text{-Hilb}(X)$ which parametrises G -clusters in X , i.e. points $\xi \in G\text{-Hilb}(X)$ correspond to 0-dimensional subschemes $Z_\xi \subset X$ such that $H^0(\mathcal{O}_{Z_\xi}) \cong \mathbb{C}[G]$, as G -representations. Typical examples of such clusters are free G -orbits. The formal definition of $G\text{-Hilb}(X)$ uses that it represents the relevant functor. Let $\tilde{X} \subset G\text{-Hilb}(X)$ be the connected component which contains the free orbits. Then there is a birational morphism (the Hilbert-Chow map) $\tilde{X} \rightarrow X/G$. Combined with the projection $X \rightarrow X/G$, this yields a universal subscheme $\mathcal{Z} \subset \tilde{X} \times X$; note that canonically $\mathcal{O}_{\mathcal{Z}} \in \text{D}^{1 \times G}(\tilde{X} \times X)$. We refer to [3] for

Theorem 7 (Bridgeland-King-Reid). *Suppose that ω_X is locally trivial in $\text{Coh}^G(X)$ and $\dim(\tilde{X} \times_{X/G} \tilde{X}) \leq \dim(X) + 1$. Then $\text{FM}_{\mathcal{O}_{\mathcal{Z}}} : \text{D}^b(\tilde{X}) \xrightarrow{\sim} \text{D}^G(X)$ is an equivalence.*

3.1 Hilbert schemes on K3 surfaces

At first consider two smooth projective varieties X and Y and the n -fold products X^n, Y^n with their natural S_n -actions. Let $P \in \text{D}^b(X \times Y)$ be the kernel

of a Fourier-Mukai transform $\mathrm{FM}_P : \mathrm{D}^b(X) \rightarrow \mathrm{D}^b(Y)$. Then, the exterior tensor product $P^{\boxtimes n} = p_1^*P \otimes \cdots \otimes p_n^*P \in \mathrm{D}^b(X^n \times Y^n)$ yields the functor $F^n = \mathrm{FM}_{P^{\boxtimes n}} : \mathrm{D}^b(X^n) \rightarrow \mathrm{D}^b(Y^n)$. Furthermore, $P^{\boxtimes n}$ has an obvious $(S_n)_\Delta$ -linearisation via permutation of tensor factors. Hence, using inflation, we get the new functor $F^{[n]} := \mathrm{FM}_{P^{\boxtimes n}}^{S_n} : \mathrm{D}^{S_n}(X^n) \rightarrow \mathrm{D}^{S_n}(Y^n)$. If we restrict to the case $X = Y$ and autoequivalences $F : \mathrm{D}^b(X) \xrightarrow{\sim} \mathrm{D}^b(X)$, we get a group homomorphism by Theorem 6

$$\mathrm{Aut}(\mathrm{D}^b(X)) \rightarrow \mathrm{Aut}^{(S_n)_\Delta}(\mathrm{D}^b(X^n)) \rightarrow \mathrm{Aut}(\mathrm{D}^{S_n}(X)), \quad \mathrm{FM}_P \mapsto \mathrm{FM}_{(P^{\boxtimes n}, \mathrm{perm})}^{S_n}.$$

From now on we need the provision $\dim(X) = \dim(Y) = 2$. It is well-known that for surfaces $\mathrm{Hilb}_n(X)$ is a crepant resolution of X^n/S_n . Furthermore, a theorem of Haiman states $\mathrm{Hilb}_n(X) \cong S_n\text{-Hilb}(X^n)$, see [7]. If we additionally assume $\omega_X \cong \mathcal{O}_X$ and $\omega_Y \cong \mathcal{O}_Y$, then we can invoke Theorem 7 in order to obtain $\mathrm{D}^b(\mathrm{Hilb}_n(X)) \cong \mathrm{D}^{S_n}(X^n)$ because in this case X^n and Y^n are symplectic manifolds and the condition $\dim(\mathrm{Hilb}_n(X) \times_{X^n/S_n} \mathrm{Hilb}_n(X)) < 1 + n \dim(X)$ of Theorem 7 is automatically fulfilled; see [3, Corollary 1.3]. However, a posteriori this inequality is true for general surfaces as the dimension of the fibre product is a local quantity which may be computed with any (e.g. affine or symplectic) model. The above homomorphism of groups of autoequivalences is now

$$\mathrm{Aut}(\mathrm{D}^b(X)) \rightarrow \mathrm{Aut}(\mathrm{D}^b(\mathrm{Hilb}_n(X))).$$

It is always injective: this is clear for $n > 2$ since the centre of S_n is trivial in this case. For $n = 2$ one can check that the sheaf $\mathcal{O}_{\Delta_X \times \Delta_X}$ with the non-canonical $(S_2)_\Delta$ -linearisation is not in the image of $\mathrm{Aut}(\mathrm{D}^b(X)) \rightarrow \mathrm{Aut}^{(S_2)_\Delta}(\mathrm{D}^b(X^2))$.

Let us introduce the shorthand $X^{[n]} := \mathrm{Hilb}_n(X)$, so that $\mathrm{D}^b(X^{[n]}) \cong \mathrm{D}^{S_n}(X^n)$ by the above and $\mathrm{Aut}(\mathrm{D}^b(X)) \hookrightarrow \mathrm{Aut}(\mathrm{D}^b(X^{[n]}))$. The above technique shows

Proposition 8. *If X and Y are two smooth projective surfaces with $\mathrm{D}^b(X) \cong \mathrm{D}^b(Y)$, then $\mathrm{D}^b(X^{[n]}) \cong \mathrm{D}^b(Y^{[n]})$. $\square$*

Remark 9. A birational isomorphism $f : X \dashrightarrow Y$ of smooth projective surfaces induces a birational map $f^{[n]} : X^{[n]} \dashrightarrow Y^{[n]}$ between their Hilbert schemes. There is a derived analogue: a Fourier-Mukai transform (resp. equivalence) $F = \mathrm{FM}_P : \mathrm{D}^b(X) \rightarrow \mathrm{D}^b(Y)$ induces a functor (resp. equivalence) $F^{[n]} = \mathrm{FM}_{P^{\boxtimes n}}^{S_n} : \mathrm{D}^b(X^{[n]}) \rightarrow \mathrm{D}^b(Y^{[n]})$. Since for the time being it is unknown whether every functor $F : \mathrm{D}^b(X) \rightarrow \mathrm{D}^b(Y)$ is of Fourier-Mukai type, we have to restrict to those (which include equivalences by [14, Theorem 2.2]).

Finally, we specialise to K3 surfaces.

Proposition 10. *Let X_1 and X_2 be two projective K3 surfaces. If there is a birational isomorphism $X_1^{[n]} \dashrightarrow X_2^{[n]}$ of their Hilbert schemes, then the derived categories are equivalent: $\mathrm{D}^b(X_1^{[n]}) \cong \mathrm{D}^b(X_2^{[n]})$.*

Proof. A birational isomorphism $f : X_1^{[n]} \dashrightarrow X_2^{[n]}$ induces an isomorphism on second cohomology, $f^* : \mathrm{H}^2(X_1^{[n]}) \xrightarrow{\sim} \mathrm{H}^2(X_2^{[n]})$, respecting the Hodge structures, because Hilbert schemes of symplectic surfaces are symplectic manifolds. From the crepant resolution $X_1^{[n]} \rightarrow X_1^n/S_n$, we find $\mathrm{H}^2(X_1) \subset \mathrm{H}^2(X_1^{[n]})$, and only the

exceptional divisor $E_1 \subset X_1^{[n]}$ is missing: $H^2(X_1^{[n]}) = H^2(X_1) \oplus \mathbb{Z} \cdot \delta_1$ with $2\delta_1 = E_1$. In particular, as $[E_1]$ is obviously an algebraic class, the transcendental sublattices coincide: $T(X_1) = T(X_1^{[n]})$. Hence, the birational isomorphism furnishes an isometry $T(X_1) \cong T(X_2)$. Orlov's derived Torelli theorem for K3 surfaces [14] then implies $D^b(X_1) \cong D^b(X_2)$. But now we apply the above result on lifting equivalences from K3 surfaces to Hilbert schemes and deduce $D^b(X_1^{[n]}) \cong D^b(X_2^{[n]})$, as claimed. $\square$

Remarks 11. (1) follows from Remark 9 and (2) from Proposition 10.

- (1) D-equivalent abelian or K3 surfaces X_1 and X_2 have D-equivalent Hilbert schemes $X_1^{[n]}$ and $X_2^{[n]}$.
- (2) Considering only birational equivalence classes of Hilbert schemes on K3 surfaces, we find that each such class is finite, since K3 surfaces have only finitely many Fourier-Mukai partners [4].
- (3) Markman [12] gives an example of non-birational Hilbert schemes $X_1^{[n]}$ and $X_2^{[n]}$ with $H^2(X_1^{[n]}) \cong H^2(X_2^{[n]})$. The above arguments still yield $D^b(X_1) \cong D^b(X_2)$ and $D^b(X_1^{[n]}) \cong D^b(X_2^{[n]})$, i.e. $X_1^{[n]}$ is D-equivalent to $X_2^{[n]}$.

3.2 Kummer surfaces

Let A be an abelian variety and consider the action of $G := \{\pm \text{id}_A\} \cong \mathbb{Z}/2\mathbb{Z}$. In order to investigate $\text{Aut}(D^G(A))$, we start with the exact sequence (see Orlov's article [15]):

$$0 \longrightarrow \mathbb{Z} \times A \times \hat{A} \xrightarrow{\eta} \text{Aut}(D^b(A)) \xrightarrow{\gamma} \text{Sp}(A \times \hat{A}) \longrightarrow 0,$$

the first morphism η maps a triple (n, a, ξ) to the autoequivalence $t_a^* \circ M_\xi[n]$, where $t_a : A \xrightarrow{\sim} A$ denotes the translation by a and $M_\xi : D^b(A) \xrightarrow{\sim} D^b(A)$ the line bundle twist with ξ . Note that shifts, translations and twists by degree 0 line bundles commute. Before turning to the second morphism γ , we set

$$\text{Sp}(A \times \hat{A}) := \left\{ \begin{pmatrix} f_1 & f_2 \\ f_3 & f_4 \end{pmatrix} \in \text{Aut}(A \times \hat{A}) : \begin{pmatrix} f_1 & f_2 \\ f_3 & f_4 \end{pmatrix} \begin{pmatrix} f_4 & -f_2 \\ -f_3 & f_1 \end{pmatrix} = \begin{pmatrix} \text{id}_A & 0 \\ 0 & \text{id}_{\hat{A}} \end{pmatrix} \right\}.$$

Now given $F \in \text{Aut}(D^b(X))$, there is a functorial way to attach an equivalence $\Phi_F : D^b(A \times \hat{A}) \xrightarrow{\sim} D^b(A \times \hat{A})$ which sends skyscraper sheaves to skyscraper sheaves. Hence Φ_F yields an automorphism $\gamma(F) : A \times \hat{A} \xrightarrow{\sim} A \times \hat{A}$ which turns out to be in $\text{Sp}(A \times \hat{A})$ (see the original [15, §2] by Orlov or [16, §4] for a slightly different presentation).

Note that G acts on $\text{Aut}(D^b(A))$ via conjugation, i.e. $(-1) \cdot F := (-1)^* \circ F \circ (-1)^*$. This induces an action on $\text{Sp}(A \times \hat{A})$, which is trivial since $\gamma((-1)^*) = -\text{id}_{A \times \hat{A}}$ is central. Taking G -invariants of Orlov's exact sequence, we get

$$0 \longrightarrow \mathbb{Z} \times A[2] \times \hat{A}[2] \xrightarrow{\eta^G} \text{Aut}(D^b(A))^G \xrightarrow{\gamma^G} \text{Sp}(A \times \hat{A}) \longrightarrow 0.$$

Here, $A[2] \subset A$ and $\hat{A}[2] \subset \hat{A}$ denote the 2-torsion subgroups; the surjectivity of γ^G uses $H^1(G, \mathbb{Z} \times A \times \hat{A}) = 0$; see [16, Proposition 4.8]. Hence, any autoequivalence $F \in \text{Aut}(D^b(A))$ differs from a G -invariant one just by translations and degree 0 line bundle twists.

Assume from now on that A is an abelian *surface*. Let X be the corresponding Kummer surface, which is a crepant resolution of A/G . A realisation is $X = G\text{-Hilb}(A)$ and we use derived McKay correspondence (Theorem 7) to infer $D^b(X) \cong D^G(A)$. From Theorem 6 we get group homomorphisms

$$\mathrm{Aut}(D^b(A))^G \xleftarrow[2:1]{for} \mathrm{Aut}^{G\Delta}(D^b(A)) \xrightarrow[2:1]{inf} \mathrm{Aut}(D^G(A))$$

In our situation both *for* and *inf* are 2:1, and *for* is surjective as $H^2(\mathbb{Z}/2\mathbb{Z}, \mathbb{C}^*) = 0$. There does not seem to be a homomorphism between the lower groups making the triangle commutative.

However, when going to cohomology (here with $\mathbb{Q}$ coefficients throughout) the roof can be completed to a diagram (refer [16, Proposition 4.14]):

$$\begin{array}{ccc} & \mathrm{Aut}(D^b(A)) & \xrightarrow{(\cdot)^H} \mathrm{Aut}(H^{2*}(A)) \\ & \nearrow for & \uparrow res \\ \mathrm{Aut}^{G\Delta}(D^b(A)) & & \\ & \searrow inf & \\ & \mathrm{Aut}(D^b(X)) & \xrightarrow{(\cdot)^H} \mathrm{Aut}(H^*(X)) \end{array}$$

Here, $(\cdot)^H : \mathrm{Aut}(D^b(A)) \rightarrow \mathrm{Aut}(H^{2*}(A))$ sends a Fourier-Mukai equivalence to the corresponding isomorphism on cohomology. Further we use that the image of $(\cdot)^H \circ inf$ lies inside the subgroup of isometries preserving the exceptional classes,

$$\mathrm{Aut}(H_{\mathrm{ex}}^*(X)) := \{\varphi \in \mathrm{Aut}(H^*(X)) : \varphi(\Lambda) = \Lambda\}$$

where $\Lambda \cong \mathbb{Q}^{16}$ is the lattice spanned by the (-2) -classes arising from the Kummer construction; the morphism $res : \varphi \mapsto \varphi|_{\Lambda}$ is then the obvious restriction.

3.3 Canonical quotients

Let X be a smooth projective variety whose canonical bundle is of finite order. Suppose that $n > 0$ is minimal with $\omega_X^n \cong \mathcal{O}_X$. Then there is an étale covering $\tilde{X} \xrightarrow{\pi} X$ of degree n with $\omega_{\tilde{X}} \cong \mathcal{O}_{\tilde{X}}$. One concrete definition is $\tilde{X} = \mathrm{Spec}(\mathcal{O}_X \oplus \omega_X \oplus \dots \oplus \omega_X^{n-1})$, see [1, §I.17]. The group $G := \mathbb{Z}/n\mathbb{Z}$ then acts freely on $\tilde{X}$ with $\tilde{X}/G = X$. Fix a generator $g \in G$ and note that a G -linearisation for some $\tilde{E} \in D^b(\tilde{X})$ is completely determined by an isomorphism $\lambda : \tilde{E} \simeq g^* \tilde{E}$ subject to $(g^{n-1})^* \lambda \circ \dots \circ g^* \lambda \circ \lambda = \mathrm{id}_{\tilde{E}}$ as G is cyclic. Here we have an equivalence $\mathrm{Coh}(X) \cong \mathrm{Coh}^G(\tilde{X})$ already on the level of abelian categories (see [13, §7]). Hence, $D^b(X) \cong D^G(\tilde{X})$ as well, a fact which also follows from the derived McKay correspondence using the trivially crepant resolution $X \xrightarrow{\mathrm{id}_X} X$. Then,

$$\mathrm{Aut}(D^b(\tilde{X}))^G \xleftarrow[n:1]{for} \mathrm{Aut}^{G\Delta}(D^b(\tilde{X})) \xrightarrow[n:1]{inf} \mathrm{Aut}(D^b(X)) = \mathrm{Aut}(D^G(\tilde{X})).$$

Bridgeland and Maciocia consider in [5] canonical quotients from the point of view of derived categories. They introduce the set of all *equivariant* equivalences

by

$$\mathrm{Aut}_{\mathrm{eq}}(\mathrm{D}^b(\tilde{X})) := \{(F, \mu) \in \mathrm{Aut}(\mathrm{D}^b(\tilde{X})) \times \mathrm{Aut}(G) : g_* \circ \tilde{F} \cong \tilde{F} \circ \mu(g)_* \forall g \in G\};$$

this is actually a group. There is an exact sequence

$$0 \longrightarrow \mathrm{Aut}(\mathrm{D}^b(\tilde{X}))^G \longrightarrow \mathrm{Aut}_{\mathrm{eq}}(\mathrm{D}^b(\tilde{X})) \longrightarrow \mathrm{Aut}(G) \longrightarrow 0$$

where the latter morphism maps $(\tilde{F}, \mu) \mapsto \mu$ and $\mathrm{Aut}(\mathrm{D}^b(\tilde{X}))^G$ is by definition the group of all equivariant equivalences with $\mu = \mathrm{id}_G$. Note that $G \cong \mathbb{Z}/n\mathbb{Z}$ implies $\mathrm{Aut}(G) \cong \mathbb{Z}/\varphi(n)\mathbb{Z}$. Furthermore, we have a subgroup $G \hookrightarrow \mathrm{Aut}(\mathrm{D}^b(\tilde{X}))^G$, $g \mapsto g_* = (g^{-1})^*$, or also $G \hookrightarrow \mathrm{Aut}_{\mathrm{eq}}(\mathrm{D}^b(\tilde{X}))$, $g \mapsto (g_*, \mathrm{id}_G)$. The latter is a normal divisor in view of $(\tilde{F}, \mu)^{-1} \circ (g_*, \mathrm{id}_G) \circ (\tilde{F}, \mu) = (\tilde{F}^{-1}, \mu^{-1}) \circ (g_* \circ \tilde{F}, \mu) = (\mu(g)_*, \mathrm{id}_G)$. By [5, Theorem 4.5] every equivalence $F \in \mathrm{Aut}(\mathrm{D}^b(X))$ has an equivariant lift $\tilde{F} \in \mathrm{Aut}(\mathrm{D}^b(\tilde{X}))$, i.e. $\pi_* \circ \tilde{F} \cong F \circ \pi_*$ and $\pi^* \circ F \cong \tilde{F} \circ \pi^*$. If $\tilde{F}_1$ and $\tilde{F}_2$ are two lifts of F , then $\tilde{F}_2^{-1} \circ \tilde{F}_1$ is a lift of $\mathrm{id}_{\mathrm{D}^b(X)}$ and thus $\tilde{F}_2^{-1} \circ \tilde{F}_1 \cong g_*$ for a $g \in G$ ([5, Lemma 4.3(a)]). Thus the equivariant lift $\tilde{F} \in \mathrm{Aut}_{\mathrm{eq}}(\mathrm{D}^b(\tilde{X}))$ is unique up to the action of G and we get a group homomorphism $\mathrm{lift} : \mathrm{Aut}(\mathrm{D}^b(X)) \rightarrow \mathrm{Aut}_{\mathrm{eq}}(\mathrm{D}^b(\tilde{X}))/G$. [5, Lemma 4.3(b)] states that if $F, F' \in \mathrm{Aut}(\mathrm{D}^b(X))$ both lift to $\tilde{F} \in \mathrm{Aut}_{\mathrm{eq}}(\mathrm{D}^b(\tilde{X}))$, then they differ by a line bundle twist: $F \cong F' \circ \mathcal{M}_{\omega_X^i}$ with $0 \leq i < n$. Thus lift is $n : 1$, and we propose the following commutative pentagram

$$\begin{array}{ccc} & \mathrm{Aut}^{G\Delta}(\mathrm{D}^b(\tilde{X})) & \\ \textit{for} \swarrow \scriptstyle n:1 & & \searrow \textit{inf} \scriptstyle n:1 \\ \mathrm{Aut}(\mathrm{D}^b(\tilde{X}))^G & & \mathrm{Aut}(\mathrm{D}^b(X)) = \mathrm{Aut}(\mathrm{D}^G(\tilde{X})) \\ \downarrow \scriptstyle 1:\varphi(n) & & \downarrow \scriptstyle n:1 \textit{ lift} \\ \mathrm{Aut}_{\mathrm{eq}}(\mathrm{D}^b(\tilde{X})) & \xrightarrow{\scriptstyle n:1} & \mathrm{Aut}_{\mathrm{eq}}(\mathrm{D}^b(\tilde{X}))/G \end{array}$$

The commutativity of this diagram boils down to the following question: given a kernel $(P, \rho) \in \mathrm{Aut}^{G\Delta}(\mathrm{D}^b(\tilde{X}))$, is $\mathrm{FM}_P : \mathrm{D}^b(\tilde{X}) \xrightarrow{\sim} \mathrm{D}^b(\tilde{X})$ a lift of $\mathrm{FM}_{(P, \rho)}^G : \mathrm{D}^G(\tilde{X}) \xrightarrow{\sim} \mathrm{D}^G(\tilde{X})$ (where we identify $\mathrm{D}^G(\tilde{X}) \cong \mathrm{D}^b(X)$)? However, this is clear from $\pi_* : \mathrm{D}^b(\tilde{X}) \rightarrow \mathrm{D}^G(\tilde{X})$, $E \mapsto \mathrm{inf}(E)$ and $\pi^* : \mathrm{D}^G(\tilde{X}) \rightarrow \mathrm{D}^b(\tilde{X})$, $(F, \lambda) \mapsto [F, \lambda]^G$.

Example 12. In the case of a double covering (e.g. X an Enriques surface), we have $n = 2$ and hence $\mathrm{Aut}(\mathrm{D}^b(\tilde{X}))^G = \mathrm{Aut}_{\mathrm{eq}}(\mathrm{D}^b(\tilde{X}))$. Then the diagram looks like

$$\begin{array}{ccc} & \mathrm{Aut}^{G\Delta}(\mathrm{D}^b(\tilde{X})) & \\ & \swarrow & \searrow \textit{inf} \\ \mathrm{Aut}(\mathrm{D}^b(\tilde{X}))^G/G & \xleftarrow{\textit{lift}} & \mathrm{Aut}(\mathrm{D}^b(X)) \end{array}$$

References

- [1] W. Barth, C. Peters, A. Van de Ven, *Compact complex surfaces*, Springer-Verlag, Berlin (1984).
- [2] J. Bernstein, V. Lunts, *Equivariant sheaves and functors*, Springer-Verlag Berlin, LNM 1578 (1994).
- [3] T. Bridgeland, A. King, M. Reid, *The McKay correspondence as an equivalence of derived categories*, J. Am. Math. Soc. 14 (2001), 535–554, also [math.AG/990802](#).
- [4] T. Bridgeland, A. Maciocia, *Complex surfaces with equivalent derived categories*, Math. Z. 236 (2001), 677–697.
- [5] T. Bridgeland, A. Maciocia, *Fourier-Mukai transforms for quotient varieties*, [math.AG/9811101](#).
- [6] C.W.C. Curtis, I. Reiner, *Representation theory of finite groups and associative algebras*, Interscience Publishers, New York-London (1962).
- [7] M. Haiman, *Hilbert schemes, polygraphs and the Macdonald positivity conjecture*, J. Amer. Math. Soc. 14 (2001), 941–1006, also [math.AG/0010246](#).
- [8] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, Berlin (1967).
- [9] Y. Ito, I. Nakamura, *McKay correspondence and Hilbert schemes*, Proc. Japan Acad. Ser. A Math. Sciences 72 (1996), 135–138.
- [10] Y. Kawamata, *D-equivalence and K-equivalence*, J. Diff. Geom. 61 (2002), 147–171, also [math.AG/0210439](#).
- [11] Y. Kawamata, *Equivalences of derived categories of sheaves on smooth stacks*, Amer. J. Math. 126 (2004), 1057–1083, also [math.AG/0205287](#).
- [12] E. Markman, *Brill-Noether duality for moduli sheaves on a K3*, J. Algebraic Geom. 10 (2001), 623–694, also [math.AG/9901072](#).
- [13] D. Mumford, *Geometric Invariant Theory*, Springer-Verlag, Berlin (1965).
- [14] D.O. Orlov, *Equivalences of derived categories and K3-surfaces*, J. Math. Sci. 84 (1997), 1361–1381, also [math.AG/9606006](#).
- [15] D.O. Orlov, *Derived categories of coherent sheaves on abelian varieties and equivalences between them*, Izv. Math. 66 (2002), 569–594, also [math.AG/9712017](#).
- [16] D. Ploog, *Groups of autoequivalences of derived categories of smooth projective varieties*, Ph-D thesis Berlin 2005.
- [17] J.P. Serre, *Galois cohomology*, Springer-Verlag, Berlin (1997).